

WELCOME

TO



This is an Education Platform

We Provide PDF Notes for Pharmacy Students

Web Site <http://www.fdspharmacy.in/>

You tube <https://www.youtube.com/c/FDSpharmacy>

Telegram <https://t.me/Fdspharmacy>

App <https://play.google.com/store/apps/details?id=com.FDSPharmacyMedia.FDSPharmacy>

E-mail fdsparmacyinfo@gmail.com

Bachelor of Pharmacy Communication skills

NOTES

✓ Unit 1 **All Unit**
✓ Unit 2 **in**
✓ Unit 3 **One PDF**
✓ Unit 4
✓ Unit 5

Visit our Website
WWW.fdspharmacy.in



Bachelor of Pharmacy Human Anatomy and Physiology I

NOTES

✓ Unit 1 **All Unit**
✓ Unit 2 **in**
✓ Unit 3 **One PDF**
✓ Unit 4
✓ Unit 5

Visit our Website
WWW.fdspharmacy.in



✓ Unit 1 **All Unit**
✓ Unit 2 **in**
✓ Unit 3 **One PDF**
✓ Unit 4
✓ Unit 5

Visit our Website
WWW.fdspharmacy.in



Bachelor of Pharmacy Pharmaceutical Inorganic Chemistry

NOTES

✓ Unit 1 **All Unit**
✓ Unit 2 **in**
✓ Unit 3 **One PDF**
✓ Unit 4
✓ Unit 5

Visit our Website
WWW.fdspharmacy.in



Bachelor of Pharmacy Remedial Biology

NOTES

✓ Unit 1 **All Unit**
✓ Unit 2 **in**
✓ Unit 3 **One PDF**
✓ Unit 4
✓ Unit 5

Visit our Website
WWW.fdspharmacy.in



Bachelor of Pharmacy Pharmaceutics I

NOTES

✓ Unit 1 **All Unit**
✓ Unit 2 **in**
✓ Unit 3 **One PDF**
✓ Unit 4
✓ Unit 5

Visit our Website
WWW.fdspharmacy.in





D.Pharma B.Pharma

- 👉 PDF Notes
- 👉 Practical Manual
- 👉 Important Questions
- 👉 Assignment etc



Download Now



www.fdpharmacy.in

REMEDIAL BIOLOGY

UNIT 5

TOPIC :

- **Cell- The unit of life**

Structure and functions of cell and cell organelles. Cell division



Cell

Cell is the basic structural & functional unit of all living organism. All the living organisms can be categorised into

- I. Unicellular organisms eg. Protozoa, yeast, and bacteria
- II. Multicellular organisms eg. Plants & humans.
- III. A human cell consist 100 trillion of cells
 - Size of the cell is 10 μm
 - Mass of the cell is 1 nano gram

Cell are two Types

- ❖ **Prokaryotic Cells** : These cells do not have well developed nucleus, but organells are present.
Eg. Archaebacteria,
- ❖ **Eukaryotic Cells** : These cells have well developed nucleus & other organells.
Eg, plants, animals


Difference Between Plant Cell and Animal Cell

Feature	Plant Cell	Animal Cell
1. Cell Wall	Present (made of cellulose); provides rigidity	Absent
2. Shape	Usually rectangular or fixed shape	Usually round, irregular, or flexible shape
3. Vacuole	One large central vacuole filled with cell sap	Many small vacuoles, or may be absent
4. Plastids	Present (e.g., chloroplasts for photosynthesis)	Absent
5. Centrosome	No distinct centrosome, but has microtubule-organizing regions	Centrosome with centrioles is present
6. Lysosomes	Rare or absent	Present (more prominent)
7. Energy Production	Uses mitochondria and chloroplasts	Uses only mitochondria
8. Nutrition	Mostly autotrophic (photosynthesis)	Heterotrophic (ingest food)
9. Cytokinesis (Cell Division)	Occurs by cell plate formation	Occurs by cleavage furrow
10. Cilia and Flagella	Rare in most plant cells	May be present (e.g., sperm cells)
11. Storage of Carbohydrates	Stored as starch	Stored as glycogen
12. Peroxisomes	Present	Present

Structure of cell

A cell Comprises of the following three basic components

- Cell membrane (or Plasma membrane)
- Cytoplasm
- Nucleus



1. Cell Membrane (or Plasma membrane)

- The cell membrane is a selectively permeable biological membrane which allow only certain substance to pass through it.
- It acts as a barrier between the inner and outer surface of the cell.
- It also known as Plasma lemma.

Properties of cell membrane

- ❑ It guards the cell against the external environment.
- ❑ It allows cellular transport between cells.
- ❑ It helps in recognising a cell.
- ❑ It divides the cell into smaller section having different structure & function.
- ❑ It allows the fusion of cell membrane with other membrane of cell organellst through special point of attachment.
- ❑ It regulates the transfer of certain specific molecules through the cell membrane e.g. Gap junction
- ❑ It enables motility of the cellular organells.

Structure (Fluid mosaic model)

- The fluid mosaic model is the most accepted one
- The fluid mosaic model of the plasma membrane was given by S.J Singer & G.L Nicholson

Function of cell

- ✓ **Protection of cells** : The receptors present on the cell membrane allow chemical messages to pass between the cell and the system, and thus provides protection to cells.
- ✓ **Separation from surrounding** : In an animal cell, it separates the internal environment from that of the external.
- ✓ **Tissue formation** : Groups of similar cells together form tissues.
- ✓ **Contribution to enzyme acting** : It controls the activity of enzymes which further participate in metabolism and immune system.
- ✓ **Particle Transportation** : This occurs by various mechanisms like, carries proteins and ion pumps.

2. Cytoplasm

- The cell consists of cytoplasm enclosed within the plasma membrane
- The cytoplasm includes various organelles present inside a cell and therefore these organelles are also referred as a cytoplasmic organelles
- Basically the cytoplasm is the fluid in which cell organelles remain suspended (mitochondria, Golgi body, nucleus, etc)

Structure of cytoplasm

The cytoplasm is divided into three parts Namely

1. Cytosol
2. Cytoplasmic organelles
3. Cytoplasmic Inclusion

② **Cytosol** : It is the fluid present inside the cell. The fluid bathes the cytoplasmic organelles

Characterstic features of cytosol include.

- ♦ Majority of the cell metabolism takes placem within the cytosol.
- ♦ They also act as ribosomes and intracellular receptors.

② **Cytoplasmic organelles** : Cell consists the organelles (via Golgi body, mitochondra, etc) have some specific characteristics that help in the maintenanceof life.

② **Cytoplasmic Inclusion** : certain bodies are present within a cell but are unable to perform their own metabolic process. These Structure also lac membrane binding them.

Function of cytoplasm

- It provide the site for cell growth and expansion
- It provide space for organelles tp remain suspended in the medium
- It also helps in the movement of several elements within the cell
- Various biochemical reactions, importance for the maintenance of life, occur in the cytoplasm
- It has enzymes which help in the breaking the large macromolecule into smaller ones so that other organelles can use them easily

3. Nucleus

- ✚ Nucleus is the spherical organelle that consists of a double layer membrane build up of phospholipids & protein.
- ✚ It is also known as the control cell.
- ✚ It consists of genetic material of the cell.

Structure of Nucleus

The Nucleus structure is composed of the following components.

- ❖ **Nuclear Envelope** : A large number of small circular or octagonal nuclear pores are present on the nuclear envelope. These nuclear pores regulate the movement of macromolecules from the nucleus to the cytoplasm and vice-versa.
- ❖ **Nucleoplasm** : The nuclear cavity of the nucleus is filled up by one of the specialised forms of protoplasm, known as nucleoplasm.
- ❖ **Chromatin material** : The chromatic material saturated with DNA, transmits the coded structural and functional information between the cell and organism.
- ❖ **Interphase chromatin** : During the interphase (the period of the cell cycle where the cell is not dividing) Two types of chromatin can be formed
 1. **Euchromatin** : It includes the active DNA.
 2. **Heterochromatin** : It mostly includes the inactive DNA.
- ❖ **Nucleolus** : The nucleus of the cell has a nucleolus also known as nucleoplasm.

It is a non-membrane bound structure containing protein and nucleic acids.

❖ **Chromosome** : The word Chromosome has been derived from the Greek word

Chroma means
colour
Soma
means *body*

A chromosome can be defined as DNA-bound protein (histone protein) that carry genetic information from generation to other

Function Of Nucleus

- ❖ It consists most of the physiological activities of the cell
- ❖ It functions to encode all the genetic information that carries the hereditary characteristics in living organisms
- ❖ It has direct involvement in reproduction (cell division)

Other Organelles

➤ Mitochondria

- Most of the eukaryotic cells contain a membrane-bound cell organells called mitochondria.
- Diameter ranging from $0.5\mu\text{m}$ – $10.\mu\text{m}$.
- Mitochondria are also know as the power house of the cell.
- Because they generate Adenosine Triphosphate (ATP) used as a source of chemical.
- Mitochondria are the chief source of energy in the cell.

Structure of mitochondria

Mitochondria are double membranous cell organelle having the following parts

1. Outer membrane : This membrane contain transport protein.

- which are helpful in transporting ions and atoms across the organelle.

2. Inner Menbrane : This membrane is involved in the formation of cristae . Cristae are

finger like projection in the matrix bearing small particle called F_1

particle or oxysomes.

Enzymes of the respiratory chain are present in cristae.

3. Mitochondrial matrix : matrix of the mitochondria is formed by fluid present within it

- This fluid contains enzymes involved in kebe's cycle.
- To oxidise fatty acids and pyruvic acid.

Function of mitochondria

- Regulates plasma membrane potential and cellular anabolism &catabolism.
- Synthesis of steroids.
- Aerobic Respiration to create energy for the functioning of cell.
- And The Main function of mitochondria , which are organelles also known as the power house of the cell because it is produce energy.

➤ Golgi bodies

→ The golgi body comprises of folded and compressed membranous sacs piled up each other. All the cell contain golgi body however it is large in those cells facilitate the synthesis and export of protein via, endoplasmic reticulum to the golgi complex.

Structure of golgi body

→ The golgi body consists of membrane bounded stacks called cisternae each cisterna consist of flattened membrane disks that carry golgi enzymes.

Function of Golgi bodies

Golgi body is involved in the synthesis of a different number of macromolecule

- **Modifying, sorting, and packing :** The golgi complex plays an important role in sorting, modifying and packing of substance for use with in the cell secretion.
- **Modification of protein :** proteins delivered by the endoplasmic recticulum are modified by the golgi body.
- **Transportation of substance :** enzymes present in the cisternae bond to the carbohydrates and phosphate group by the process of glycosylation and phosphorylation in order to modify substance by the transport of substance.
- **Formation of spindle :** The golgi body is involved in the formulation of meiotic and mitotic spindles.
- **Excretion of water :** These organelles help in the excretion excess amount of water.

➤ Lysosomes

→ In 1949 a belgian cytologist Christian de duve had discovered the cell organelle 'lysosomes' 'The word lysosome derived from greek word

- **Lysis** means to **separate**
- **Some** means **body**

→ They are generally known as suicide-sacs or suicide bags.

Structure of lysosome

- Some non-green plant cells possess lysosomes.
- They are also called suicide bags as they cause intracellular digestion (deconstruct the carbohydrates, protein, DNA, RNA etc) and thus destroy the cell.

Characteristic feature of lysosome

- They are present in high number in a single cell.
- They possess nearly 40 different enzymes.
- They are referred as suicidal bags since the enzymes present in them can digest the whole cell in case of damage or death of the cell.

Function of lysosome

- Provide defence against invading microorganisms via circulating macrophages.
- Helps dissolve blood clots.
- Cause the sperm to penetrate the ovum.
- Decompose the mature RBC and dead cells.
- Help in reabsorption of water and small molecule in kidney and urinary bladder.

➤ **Microsomes / Endoplasmic reticulum**

- Microsomes are a heterogeneous set of vesicles having diameter 20- 200 nm consisting of endoplasmic reticulum.
- These vesicles are isolated by differential centrifugation and form three structural features.
i.e, rough vesicles smooth vesicles ribosomes.

Structure of microsomes / endoplasmic reticulum

- ⦿ **Rough endoplasmic reticulum (RER)** : Small particles of RNA known as ribosomes are found in close association with the membrane of the endoplasmic reticulum which gives a rough appearance to the membrane.
- ⦿ **Smooth endoplasmic reticulum (SER)** : The membrane of SER does not contain ribosomes and help in the metabolism of carbohydrates and lipid they are involved in the synthesis of steroid hormones in adrenals and testes.
- ⦿ **Sarcoplasmic reticulum (SR)** : Most of the muscles contain a unique type of smooth ER called sarcoplasmic reticulum.

Function of Microsome / endoplasmic reticulum

- It acts as a structural framework of the cytoplasm.
- It exchanges material with the cytosol by active transport, osmosis, and diffusion.
- Protein synthesised by its surface-bound ribosomes are collected and stored by RER.
- It has different metabolic enzymes which helps in the synthesis of phospholipid, cholesterol

➤ Ribosomes

→ Ribosomes are cell organelles where protein synthesis takes place. They are abundant in living cells. They are either found as free particles (in prokaryotic and eukaryotic cells) or attached to membrane (as in eukaryotic cells).

Structure of Ribosomes

- Ribosomes are ribonucleoproteins i.e. made from RNA Complex and protein. It is divided into smaller parts called subunits.
- Which are made up of both protein and ribonucleic acid components. Smaller sub unit & Larger sub unit.s

Function of Ribosomes

- ❑ The primary Function of ribosomes is protein synthesis.
- ❑ The information carried in the genetic code is converted into protein molecule within the ribosomes.

Function of Cell

1. cell growth & Metabolism :

- Cell grow either by increasing its number or in size
- Metabolism is the chemical reaction that occurs in our body cell its change food in energy
- During metabolism, various anabolic & catabolic process occur

Catabolic process : Breakdown of complex molecule occurs within the cells to produce energy

Anabolic process : Cells utilise energy to synthesis complex biomolecule and perfrom other biological function

2. Cell Division : It involve the splitting of a single cell. Mother cell splits into two daughter Cells this result in the growth of multicellular organisms. During the cell division, DNA also synthesised and the process is known as replication.

3. Protein Synthesis : It occurs within the cell, by two major steps
Transcription & Translation

4. Molecule Transport : The cell must be able to acquire nutrients and

other molecule to survive

Cell Division

- **Cell Division** is the process by which a parent cell divides into two or more daughter cells.
- It is essential for growth, repair, reproduction, and development in living organisms.
- There are two main types of cell division
 - Mitosis
 - Meiosis

Mitosis

- Mitosis is the process by which a single cell divides to form two identical daughter cells. It is used for growth, healing, and replacing old cells in the body.
- Also known as (M-phase or mitotic phase (Equational division)).
- It occurs in somatic cell in which already existing parent cell divides into two identical daughter cells.
- In this, the no. of chromosomes in daughter cell is same as the no. of chromosomes in parent cell
- It occurs into two parts

1. Karyokinesis : It is a process by which the cell nucleus divides into two daughter nuclei.

Prophase

- This phase begins with initiation of condensation of the chromosomal material ie. chromatin
- Centrioles (centrosome) move towards the opposite poles of the cell.

- Organelles like golgi complex, ER, Nucleolus, and Nuclear envelope disappear at the end of prophase

Metaphase

- This phase begins with complete disintegration of nucleat envelope.
- chromosomal condensation completed and two sister chromatids are holds together by centromere
- Kinetochores (small disc shaped) at the surface of centron.
- Spindle fible attache to kinetochores and chromosor are moved to spindle equator and get alligned.

Anaphase

- Begins with splitting of each chromosomes.
- Separation of chromatids and move towards Opposite poles.

Telophase

- last stage of mitosis and in this, cromosomes loose individuality (decondensation) and chromatin material is collected as a mass at two poles.
- Nuclear membrane reappear and two nudear formed
- other organelles reappears.

2. Cytokinesis

It is the division of parent cell cytoplasm after nucleus division to form two daughter cells

Meiosis

- It occurs in gamete cells (reproductive cells).
- In this, offsprings are produced by the fusion of a Female gamete and a male gamete
- It occurs in two stages :-

i) Meiosis -I

Prophase-I
Leptotene
Zygotene
Pachytene
Diplotene
Diakinesis
Metaphase-I
Anaphase-I
Telophase -I

ii) Meiosis - II

Prophase-II
Metaphase-II
Anaphase-II
Telophase-II

i) Meiosis-I→ It is divided into four phases.

Prophase-I : It is longer and more complex,

It is further sub-divided into five phases-

- a) leptotene phase → Condensation of chromosomes starts.
- b) Zygotene phase → pairing of chromosomes, and the paired chromosome referred as Homologous chromosomes process is known as Synapsis) and formed synaptonemal complex
- c) Pachytene phase → exchange of genetic material b/w two homologous chromosomes i.e. crossing over
- d) Diplotene phase → dissolution of synaptonemal complex and formed x-shaped structure termed as chiasmata)
- e) Diakinesis phase → Terminalisation of chiasmata, chromosome condensation completed, mitotic spindles are assembled, Nucleolus and nuclear envelope disappear.

Metaphase - I : chromosomes align at the equatorial plate, microtubules are seen attaching to the pair.

Anaphase - I : Homologous chromosomes separated.

Telophase - I : Nuclear membranes and nucleolus reappear. This telophase followed by cytokinesis and formed two haploid (gametes) cells.

ii) Meiosis - II : It is similar as normal mitosis. It also comprises of four stages

Prophase-II : condensation of chromosomes, formation of Spindles and disappearance of nucleus.

Metaphase-II : equatorial alignment of Chromosomes.

Anaphase-II : separation of chromatides and move towards opposite Poles

Telophase-II : chromosomes loose individually and chromatin material is called as mass of two poles. Nuclear membranes reappear from organelles

