WELCOME

TO



This is an Education Platform

We provide Free PDF Notes and Videos Classes for Pharmacy Students

Web Site http://www.fdspharmacy.in/

You tube https://www.youtube.com/channel/UC77iEsiuZolU4pB8WAJIR5Q

What app https://chat.whatsapp.com/IzSgXtFEvhS4LN5xhUgq5z

Telegram https://t.me/+cvxm17xSloA4MjVl

Face book https://www.facebook.com/Fdspharmacy-105764311994440/

E-mail fdspharmacyinfo@gmail.com

Diploma in Pharmacy 2nd Year
Pharmacology
Important Questions
Chapter 3 : Drugs Acting on Eye

Questions	Page No
Q1. What is miotics ? Write the classification, pharmacological action, Indication, contraindications, dose, of miotics.	3
Q2. Explain the Mydriatics ? Write the classification, pharmacological action, Indication, contraindications, dose, of Mydriatics.	4
Q3. Write the note on Glaucoma.	5

FDSPharmacy
Learn and Educate

Chapter 3
Drugs Acting on Eye
IMPORTANT Questions

Q1. What is miotics ? Write the classification, pharmacological action, Indication, contraindications, dose, of miotics.

Ans.

Miotics

- The drugs are used to produce miosis / constriction of pupils are called Miotics.

Classification

1. Muscarinic Agents : Pilocarpine , carbachol and other cholinergic drugs .
2. α adrenergic blockers(relaxation of dilator muscle) : Tolazoline and other alpha blocker . pupil get constricted when iris dilated .

Indications

- These drugs are used to treat glaucoma .
- To control mydratic effect.
- To prevent or break adhesion (attachment) between iris and lens.

Contraindications

- These are contraindicated during pregnancy.
- These are contraindicated during lactation.
- These are contraindicated during in children because its safety is not clear.
- These are contraindicated in patients are above 65 years.
- These are contraindicated in patients are suffering from cataract.

Dose

- Pilocarpine : 1-4% pilocarpine nitrate in eye drops.
- Tolazoline ; 1-2mg/kg intravenously over 10 minutes.

Q2. Explain the Mydriatics ? Write the classification, pharmacological action, Indication, contraindications, dose, of Mydriatics.

Ans.

Mydriatics

- The drug are used to dilate the pupil to perform easier and detailed examination.

Classification

1. Adrenergic agonist : Adrinaline , Phenylephrine.
2. Cholinergic Antagonist : Tropicamide , Atropine , cyclopentolet.

Indications

- These are used to dilate the pupil.
- These are used for examination of retina and optic nerves.
- These are used for diagnosis of iris problems.
- These are used to determine refractive error (it is a disease in which eye can not focus the image clearly and result in blurred vision.

Contraindications

- It is contraindicated in hypertension.
- In Glaucoma.
- In intraocular pressure.

Doses

- Tropicamide : 0.5% solution dilate for 30 minutes nearly.
- Atropine : 1% solution 7-12 days
- Cyclopentolate : 1-2 % 24 hours

Q3. Write the note on Glaucoma.

Ans.

Glaucoma

- → Glaucoma is a ocular (eye) Disease in which intraocular pressure is increased and damage the optic nerves which causes visual loss.
- → When drainage canal of aqueous humorous get blocked , the intraocular pressure increases.

There are two types of glaucoma :

- Open Angle Glaucoma
- Narrow / Closed Angle Glaucoma

Classification

1. Osmotic Agent : Mannitol, Glycerol
2. β- Adrenergic blockers : Timolol, Betaxolol, Brimonidine
3. Miotics : Pilocarpine,

Indication

- For the treatment of glaucoma
- Reduce Intraocular Pressure
- Eye pain etc

Dose

- Timolol 0.23 – 0.5 % one drop B.D
- Brimonidine 0.2% B.D

Contraindication

- Peptic Ulcer
- Hyyotension
- Epilepsy etc

FDSPharmacy
Learn and Educate